


METEOROLOGY

Evaluation of the impact of the space component of the Global Observing System through Observing System Experiments



This article appeared in the Meteorology section of ECMWF Newsletter No. 113 – Autumn 2007, pp. 16-28.

Evaluation of the impact of the space component of the Global Observing System through Observing System Experiments

Graeme Kelly, Jean-Noël Thépaut

As many more satellite instruments (both active and passive) became operational the challenge was, and still is, how to assimilate these space measurements together with all the conventional measurements. Figure 1 shows most of the current and near future satellite sensors used or soon to be used. Since 2001 the number of satellite sensors being used has increased by a factor of four and now is in excess of forty sensors. A recent example of the number of observations, mostly satellite radiances, used in the operational data assimilation at ECMWF during a 12-hour period is given in Table 1. Complex data processing techniques are employed to assimilate all these observations in a timely manner

A study, sponsored by EUMETSAT, has been carried out to evaluate the impact of the space component of the Global Observing System through Observing System Experiments. In this study the relative contributions of the various space observing systems have been assessed within the context of the ECMWF data assimilation system. It is found that all the space based sensors generally contribute in a positive way to the overall improvement of the ECMWF forecast system.




Figure 1 Number of satellite sensors that are or will be soon assimilated in the ECMWF operational data assimilation.

	Screening		Used in Analysis	
SYNOP	421,000	0.43%	64,000	1.94%
Aircraft reports	519,000	0.53%	247,000	7.53%
DRIBU	24,000	0.02%	6,000	0.18%
TEMP	152,000	0.16%	75,000	2.28%
PILOT	119,000	0.12%	57,000	1.75%
AMVs	4,272,000	4.37%	131,000	3.99%
Radiance data	91,786,000	93.91%	2,508,000	76.46%
Scatterometer winds	274,000	0.28%	118,000	3.61%
GPS radio occultation	167,000	0.17%	73,000	2.24%
TOTAL	97,734,000	100.00%	3,280,000	100.00%

Table 1 Observation data count for one 12 hour 4D-Var cycle for 0900–2100 UTC on 24 April 2007. ‘Screening’ refers to the actual amount of data presented to 4D-Var and the ‘Used in Analysis’ indicates the amount of data used during the analysis minimization.

Requirement for Observing System Studies

At its meeting at ECMWF on 3 May 2003, the EUCOS Scientific Advisory Team discussed the need to investigate the interdependencies between the space-based and terrestrial components of the observing system. It was suggested that such an investigation could be based on a set of carefully designed Observing System Experiments (OSEs). These studies would be designed to provide guidance on the future development of the terrestrial observing system in view of the increasing capabilities of the satellite observing systems provided by the meteorological space agencies.

In recent years, several NWP centres have demonstrated substantial benefit from the assimilation of, for example, ATOVS radiances and scatterometer winds (referred to hereafter as SCAT). Since 2003 data has become available from second-generation radiometers (AIRS on Aqua in 2003 and IASI on MetOp in 2007) providing significantly enhanced temperature and humidity sounding capabilities – to be followed (in the five to ten year time frame) by similar instruments on the operational NPOESS series of satellites.

It was agreed that, as far as EUCOS is concerned, the primary issues were:

- What are the relative contributions of various components of the terrestrial observing system within the current overall composite observing system?
- How should the terrestrial systems evolve over the next five to ten years and beyond to complement the projected evolution of the space-based observing systems?

This led to a proposal from *Andersson et al.* (2004) to carry out a set of OSEs specifically designed to evaluate the role of the terrestrial component of the Global Observing System.

Following a number of discussions between EUMETSAT, ECMWF and EUCOS, it was agreed that specific OSEs dedicated to examining the various contributions of the different components of the space observing system were necessary to complement the original proposal about the terrestrial components. Taking this approach would provide a comprehensive assessment of the space/terrestrial links. It was also agreed that the robustness of this combined assessment would be strengthened by the adoption of similar strategies for experimentation and validation of the two studies.

These studies also take onboard one of the outcomes of the Third WMO Workshop on the Impact of Various Observing Systems on NWP. This suggested that, due to a large degree of redundancy of the Global Observing System (GOS), performing impact studies by removing one element of the GOS can show very limited impact and does not necessarily highlight the intrinsic benefit of the element in question. It was therefore decided that the scenarios in which the contributions of different elements of the GOS are investigated would be based on adding datasets or combination of datasets to a reference.

This article will deal with the relative contributions of the various space observing systems which have been assessed within the context of the ECMWF data assimilation system. The results of the complementary study concerning the contributions from various terrestrial observing systems are summarised in Box A.

Terrestrial observing system studies (EUCOS)

A

The set of terrestrial observing system studies coordinated by EUCOS has been completed following the guidelines indicated in *Andersson et al.* (2004). These impact studies aimed to examine the various components of the terrestrial observing system, in the presence of the current satellite-based observing system. The experiments have been run using the same first winter and summer period used for the space observing system studies with the identical assimilation setup to enable a direct comparison with the space studies. The total number of cases remains probably too short to provide statistical robustness to the findings (especially over small areas such as Europe), but it is reassuring that the impact of the various components of the terrestrial observing system remains similar to the first order between the two assessed periods.

The main findings of the winter impact studies indicate a large impact of the radiosondes (wind and temperature) and aircrafts (wind and temperature), a marginal impact of radiosonde humidity information, and a neutral impact from the wind profilers. Sole wind or temperature information from radiosondes is not sufficient to impact noticeably on the forecast skill. In contrast, coupled temperature/wind information from radiosondes seems to provide a large and significant improvement in the forecasts well into the medium-range. The experiments demonstrate that observations from aircraft and radiosondes are complementary: each observing system improves the forecast skill even in the presence of the other.

The summer impact studies confirm most of the findings from the winter experiments, although the impact of the various assessed components of the GOS is smaller, both in absolute and relative terms (*Thépaut & Kelly, 2007*).

Past OSEs (1996–2003) with satellite data

Three sets of OSEs were run at ECMWF soon after the introduction of 3D-Var in 1996 (Kelly, 1997), after the operational implementation of 4D-Var in 2000 (Bouttier & Kelly, 2001), and later in 2003 (Kelly et al., 2004). For each set of OSEs there were four scenarios considered.

- **CONTROL:** For each set of OSEs the model cycle closest to the operational system at that time was used.
- **NOAIREP:** All aircraft measurements (wind and temperature) removed.
- **NOUPPER:** All TEMP, PILOT and PROFILER reports removed.
- **NOSAT:** All satellite data removed (the terrestrial network used in operations).

As well as conventional observations (TEMP, PILOT, PROFILER, AIREPS, SYNOP, PAOBS and BUOY reports), the 2003 OSEs included data from:

- Three AMSU-A/B and two HIRS instruments from the NOAA satellites.
- Five geostationary satellites and one polar orbiter (Terra) providing Atmospheric Motion Vectors.
- Three geostationary satellites providing clear-sky water-vapour radiances (CSRs).
- Three SSM/I instruments from the DMSP platforms.
- Seawinds instrument from Quikscat.

The results from the three sets of OSEs are summarized in Figures 2 and 3. In a nutshell, they show that satellite data has progressively become the most important data source in both hemispheres, transcending even the conventional upper-air network in the northern hemisphere in the last set of OSEs.




Figure 2 Comparison of three OSEs in the northern hemisphere for the rms error of the 500 hPa geopotential height for (a) 2003, (b) 2000 and (c) 1996.




Figure 3 Comparison of three OSEs in the southern hemisphere for the rms error of 500 hPa geopotential height for (a) 2003, (b) 2000 and (c) 1996.

Studies of the various space observing systems

We now consider the relative contributions of the various space observing systems (infrared temperature soundings, microwave temperature soundings, imagers, scatterometers, etc.) within the context of a more recent ECMWF data assimilation system.

Reference systems

We have assumed in this study that the current conventional observing system is maintained (thereafter called the *BASELINE* system), and the main focus is to evaluate how specific satellite systems contribute individually to the robustness of the GOS, in addition to this degraded observing network.

The evaluation of satellite sensors is best done in the tropics and southern hemisphere, but the quality of the *BASELINE* system (equivalent to *NOSAT* referred to earlier) is so poor outside the northern hemisphere that it was not considered suitable as a reference by itself. Instead, two special reference systems have been designed to ensure a reasonable quality of the atmospheric analyses and forecasts in the tropics and southern hemisphere. These special reference systems are:

- **AMV(REF):** *BASELINE* plus the Atmospheric Motion Vectors (AMVs).
- **AMSUA(REF):** *BASELINE* plus data from one AMSU-A instrument.

Experimental setup

The data assimilation framework used for all the OSEs corresponds to the system that was used in operations close to the time of the periods under investigation. The main characteristics of the data assimilation system are given in Box B.

The experiments were carried out for two periods, each covering a winter and summer.

- **Period 1.** Winter from 4 December 2004 until 25 January 2005 and summer from 17 July to 15 September 2005 using IFS Cy29r1 and Cy29r2 respectively. For this period, the OSEs were based on *AMV(REF)* as a reference.
- **Period 2.** Winter from 5 December 2006 to 14 February 2007 and summer from 1 June to 18 August 2006 using IFS Cy31r1. For this period, the OSEs were based on *AMSUA(REF)* as a reference.

All forecasts were run from 00 UTC.

Standard experiments

Two sets of assimilation were performed.

- **AMV(REF) as reference for Period 1.** The observational scenarios tested with AMVs as reference (i.e. *AMV(REF)*) are described in Table 2. These experiments are based on the winter and summer forming Period 1. The first ten days of each assimilation scenario are excluded from the verification to ensure a reasonable warm-up phase. No real difference in the impact was found between summer and winter so the mean scores are combined to give a sample of 89 days for each experiment. All experiments are validated using the operational analysis.
- **AMSUA(REF) as reference for Period 2.** The observational scenarios tested with one AMSU-A as reference (i.e. *AMSUA(REF)*) are described in Table 3. These experiments are based on the winter and summer forming Period 2. These experiments were delayed as long as possible in order to make use of the AMSU-A and MHS instruments from the EUMETSAT MetOp satellite. The first two weeks are excluded from the verification to ensure a reasonable warm-up phase for each assimilation scenario. For Period 2, the Variational Bias Correction for satellite radiances was operational and therefore activated during the warm-up phase of the experiments (bias correction coefficients are then kept constant for the remaining of the assimilation period). No real difference in the impact was found between summer and winter so the mean scores are combined to give a sample of 117 days for each experiment. All experiments are validated using the operational analysis.

Additional experiments

During the course of the study two additional sets of experiments using *AMV(REF)* and *AMSUA(REF)* have been carried out to specifically assess the impact of MODIS and AVHRR AMVs, the impact of various AIRS channel combinations (as a scientific preparation for the assimilation of IASI), and finally the respective contribution of clear and cloud/rain effected SSMI radiances.

The experiments using *AMV(REF)* are as follows.

- **Polar wind impact:** An experiment evaluating the impact of the MODIS winds by removing those winds from *AMV(REF)*.
- **Temperature and humidity impact:** An experiment adding AMSU-A and AMSU-B data to *AMV(REF)* to assess their impact compared to that of AIRS data.

The following additional experiments have been conducted using *AMSUA(REF)* as a reference.

- **Impact of various AIRS channel combinations:** Four experiments (summer only) adding various combinations of AIRS channels. A data denial experiment has also been run by removing AIRS data from *CONTROL*.
- **SSMI impact:** Three experiments (summer only) adding SSMI (clear sky), SSMI(rainy) and SSMI(clear sky + rainy) data to the *AMSUA(REF)*. In addition, two data denial experiment have also been run: removing SSMI(clear sky) and SSMI(rainy) data from *CONTROL*.
 - SSMI(clear sky) are those SSMI radiances considered to be not affected by cloud or rain using a regression based cloud liquid check.
 - SSMI(rainy) are those radiances that fail the previous cloud liquid test and pass the convergence test in the SSMI 1D-Var (*Bauer et al., 2002, 2005a and 2005b*).
- **Polar wind impact:** An experiment (winter only) adding each wind set to *AMSUA(REF)* to evaluate the impact of the MODIS and AVHRR winds.

Main characteristics of the data assimilation system**B**

- T511L60 forecast model resolution
- 4D-Var assimilation with a 12-hour window and the analysis inner and out loop resolutions being T95/T159L60 and T511L60 respectively
- Conventional observations currently assimilated in the system include:
 - TEMP, PILOT and PROFILER reports
 - SYNOP, SHIP, METAR and BUOY (moored and drifters) reports
 - Aircrafts (AMDAR, AIREP, ACARS) including ascent/descent reports
- Satellite observations assimilated in the system for the atmospheric analysis were at that time for the winter run:
 - Atmospheric Motion Vectors from GEO (Meteosat-5/7, GOES-9/10/12) and LEO (MODIS Terra and Aqua) platforms
 - Clear-sky water vapour radiances from GEO (Meteosat-5/8, GOES-9/10/12)
 - Level 1c infrared radiances from NOAA 14/17 (HIRS) and Aqua (AIRS)
 - Level 1c microwave radiances from NOAA-15 (AMSU-A), NOAA-16 (AMSU-A and AMSU-B), NOAA-17 (AMSU-B), Aqua (AMSU-A) and DMSP 13/14/15 (SSM/I)
 - Sea surface winds from scatterometers QuikScat and ERS-2
 - Ozone products from NOAA-16 (SBUV) and ENVISAT (SCIAMACHY).
- As this study has been spread over two years, different model cycles have been used for the two scenarios.
- **Period 1.** IFS model cycles Cy29r1 (winter) and Cy29r2 (summer) have been used, differing mainly by the inclusion of NOAA-18 level-1c radiances from AMSU-A and MHS and the blacklisting of NOAA-14 HIRS radiances that had become too noisy. *AMV(REF)* was used as a reference for Period 1.
- **Period 2.** IFS model cycle Cy31r1 has been used for both winter and summer. *AMSUA(REF)* was used as a reference for Period 2.

Experiment	Datasets
<i>BASELINE</i>	All conventional observations used in NWP (radiosonde + aircraft + profiler network + surface land data + buoy observations + ship data)
<i>AMV(REF)</i>	<i>BASELINE</i> + AMVs from GEO & MODIS
<i>AMV(REF)+HIRS</i>	<i>AMV(REF)</i> + HIRS radiances
<i>AMV(REF)+AMSUA</i>	<i>AMV(REF)</i> + AMSU-A radiances
<i>AMV(REF)+AMSUB</i>	<i>AMV(REF)</i> + AMSU-B radiances
<i>AMV(REF)+SSMI</i>	<i>AMV(REF)</i> + SSMI radiances
<i>AMV(REF)+CSRs</i>	<i>AMV(REF)</i> + CSRs (Clear Sky Radiances) from GEO
<i>AMV(REF)+AIRS</i>	<i>AMV(REF)</i> + AIRS radiances
<i>AMV(REF)+SCAT</i>	<i>AMV(REF)</i> + SCAT winds
<i>AMV(REF)+AMVs</i>	<i>AMV(REF)</i> + AMVs from GEO
<i>CONTROL</i>	Full operational system (all the observations)

Table 2 Observational scenarios tested with *AMV(REF)* (*BASELINE* plus AMVs from GEO and MODIS) as reference for Period 1.

Experiment	Datasets
<i>BASELINE</i>	All conventional observations used in NWP (radiosonde + aircraft + profiler network + surface land data + buoy observations + ship data)
<i>AMSUA(REF)</i>	<i>BASELINE</i> + AMSU-A radiances from NOAA-16
<i>AMSUA(REF)+AMVs</i>	<i>AMSUA(REF)</i> + AMVs from GEO & MODIS
<i>AMSUA(REF)+AMSUA</i>	<i>AMSUA(REF)</i> + AMSU-A radiances
<i>AMSUA(REF)+AMSUB</i>	<i>AMSUA(REF)</i> + AMSU-B radiances
<i>AMSUA(REF)+CSRs</i>	<i>AMSUA(REF)</i> + CSRs (Clear Sky Radiances) from GEO
<i>AMSUA(REF)+AIRS</i>	<i>AMSUA(REF)</i> + AIRS radiances
<i>AMSUA(REF)+SCAT</i>	<i>AMSUA(REF)</i> + SCAT winds
<i>CONTROL</i>	Full operational system (all the observations)

Table 3 Observational scenarios tested with *AMSUA(REF)* (*BASELINE* plus AMSU-A from NOAA-16) as reference for Period 2.

Assimilation assessment

In order to assess the impact of each OSE several statistical quantities have been calculated for temperature, humidity and wind at various levels. These include anomaly correlations, mean rms errors, geographical maps of rms error differences and mean rms error differences along with statistical significance. All these quantities have been computed but only a small selection is shown here; the remainder will be contained in a comprehensive report to be delivered to EUMETSAT. Most results shown are for the southern hemisphere or tropics where the impact is largest. In the northern hemisphere the impact is generally similar but reduced.

As stated above, in the southern hemisphere and tropics the *BASELINE* assimilation (terrestrial observations only) is poor and not suitable as a reference. The addition to the *BASELINE* of either AMVs (geostationary and polar) or data from one AMSU-A instrument considerably improves the quality of the assimilation with the southern hemispheric forecast skill increasing by about two days at day 4.

Figure 4 shows the mean anomaly correlation at 500 hPa for the combined summer and winter assimilation sets of Period 1. The experiments are:

- *BASELINE* (*NOSAT*), *AMV(REF)* and *CONTROL* as described in Table 2.
- *EUCOS(REF)* which uses all satellite data and a reduced terrestrial network, i.e. the GCOS Upper Air Network (*GUAN*), GCOS Surface Network (*GSN*), and the buoys' network.

Figure 5 is similar to Figure 4 but for Period 2 and using *AMSUA(REF)* in place of *AMV(REF)*. The *EUCOS* experiments have not been not run for this period.

For the northern hemisphere all experiments reach day 6 before the anomaly correlation drops to 0.75; this indicates the forecasts are of general good quality. When the anomaly correlation drops below 0.6 the forecasts are considered poor. In the southern hemisphere the *BASELINE* assimilations for both Period 1 and Period 2 are poor; their anomaly correlations reach 0.6 soon after day 5.

EUCOS(REF) is also shown for comparison with the satellite references (see Box B). With the addition of the remaining terrestrial observations the forecast improves by 8 hours at day 6 in the northern hemisphere. On the other hand, in the southern hemisphere, as expected, *EUCOS(REF)* is close to *CONTROL*.




Figure 4 Comparison of *EUCOS(REF)* and *AMV(REF)* with *BASELINE (NOSAT)* and *CONTROL* for (a) northern hemisphere (20°–90°N) and (b) southern hemisphere (20°–90°S).




Figure 5 Comparison of *AMSUA(REF)* with *BASELINE (NOSAT)* and *CONTROL* for (a) northern hemisphere (20°–90°N) and (b) southern hemisphere (20°–90°S).

Impact of sensors

To present the results of eleven sets of data assimilation experiment in a concise way is a somewhat daunting task. There are two sets of OSEs based on *AMV(REF)* and *AMSUA(REF)*. There is also a variety of variables and levels used for evaluation: 500 hPa geopotential height, relative humidity at 850, 500 and 200 hPa, and wind at 1000 and 200 hPa.

The results have been condensed into a series of bar graphs containing all experiments. Generally the sensors are ranked in order of increasing rms error for the first verified forecast range. Usually this ranking is maintained throughout the forecast, though there are some exceptions.

Generally all sensors impact in a positive way on some parameters but some sensors have a neutral or slightly negative impact on other parameters. The small negative impact, mostly noticed on the 500 hPa geopotential height parameter and when using *AMV(REF)* as a reference, may be due to the fact that the accuracy of the *AMV(REF)* temperature field is still not quite good enough to assimilate radiances that are mostly sensitive to moisture. This negative impact of some sensors is not generally found when *AMSUA(REF)* is used as a reference instead.

500 hPa geopotential height

The accuracy of the 500 hPa geopotential height forecast is an important and classical measure of forecast skill.

- **OSEs based on *AMV(REF)*.** Figure 6 shows the performance of all the OSEs as described in Tables 2 at days 2, 5 and 7. The largest impact can be seen in the southern hemisphere and is maintained throughout the full forecast range. Clearly the most important sensors are AMSU-A and AIRS followed by HIRS. All other sensors have a relatively small impact; some sensors even show a small negative impact relative to *AMV(REF)* for this particular parameter. However, other scores are improved by these sensors (this is for example the case for the CSRs which improve the humidity scores). The impact in the northern hemisphere is similar to that in the southern hemisphere but smaller in magnitude.
- **OSEs based on *AMSUA(REF)*.** The performance of all the OSE experiments as described in Table 3 at days 2, 5 and 7 is shown in Figure 7. First of all, it is worth noticing that the relative difference for 500 hPa geopotential height between *AMSUA(REF)* and *CONTROL* compared to the relative difference between and *AMV(REF)* and its *CONTROL* is smaller, and therefore gives less margin to measure quantitatively the impact of individual sensors. However, the largest sensor impacts can still be seen in the southern hemisphere and these are maintained throughout the full forecast range. Clearly the most important sensors are AIRS and the AMSU-A/B combination. All other sensors have a relatively small impact. In the northern hemisphere the impact of the sensors is similar but smaller in magnitude.




Figure 6 Impact of all sensors (based on *AMV(REF)*) on 500 hPa geopotential height for (a) southern hemisphere (20°–90°S) and (b) northern hemisphere (20°–90°N).




Figure 7 Impact of all sensors (based on *AMSUA(REF)*) on 500 hPa geopotential height for (a) southern hemisphere (20°–90°S) and (b) northern hemisphere (20°–90°N).

850 hPa relative humidity

Moisture forecasts, particularly in the tropics, tend to be less accurate than forecasts of mid-latitude geopotential height. After day 4, the moisture forecast becomes less dependent on the initial moisture conditions and the model moisture processes dominate. For this reason all the moisture validations are presented for days 1 to 3.

- **OSEs based on *AMV(REF)*.** Figure 8(a) shows the performance of all the OSEs as described in Table 2 for the tropics. SSMI is the most important sensor at day 1 but by day 3 the impact is reduced and overtaken by that of AIRS. However the gap between the *CONTROL* and the *AMV(REF)+SSMI* at day 1 is much larger than the difference between *AMV(REF)+SSMI* and *AMV(REF)* suggesting it is the combination of all sensors that is important rather than a single sensor. The impact in the northern hemisphere and southern hemisphere is similar to that in the tropics but smaller.
- **OSEs based on *AMSUA(REF)*.** Figure 8(b) shows the performance of all the OSEs as described in Table 3 for the tropics where the impacts are the largest. In this set, *AMSUA(REF)+SSMI* now includes both clear-sky and rain/cloud affected radiances and SSMI is the most important sensor for low level humidity. However there is still a gap between the *CONTROL* and *AMSUA(REF)+SSMI*, which again suggests that it is the combination of all sensors that is important for improving the moisture analysis and forecasts rather than a single sensor. The impact in the northern hemisphere and southern hemisphere is similar but smaller in magnitude than in the tropics.

1000 hPa wind

Wind forecasts in the tropics tend to be less accurate than in mid latitudes. In the tropics after day 4 the model wind forecast becomes less dependent on the initial conditions. Therefore all the wind validations presented here are for days 1, 2 and 3.

- **OSEs based on AMV(REF).** Figure 9(a) shows the performance of all the OSEs as described in Table 2. In the tropics SSMI is the most important sensor. However, at day 1 the gap between *CONTROL* and *AMV(REF)+SSMI* is much larger than the difference between *AMV(REF)+SSMI* and *AMV(REF)* suggesting again that it is the combination of all sensors that is important. The impact in the northern hemisphere and southern hemisphere is similar to that in the tropics but smaller in magnitude.
- **OSEs based on AMSUA(REF).** Figure 9(b) shows the performance all the OSEs as described in Table 3. In the tropics SSMI is the most important sensor, though SCAT winds are also important in the early part of the forecast. However the gap between the *CONTROL* and *AMSUA(REF)+SSMI* suggests it is the combination of all sensors that is important. The impact in the northern hemisphere and southern hemisphere is similar but smaller in magnitude to that in the tropics.




Figure 8 Impact of all sensors on 850 hPa relative humidity for the tropics based on (a) *AMV(REF)* and (b) *AMSUA(REF)*.




Figure 9 Impact of all sensors on 1000 hPa vector wind for the tropics based on (a) *AMV(REF)* and (b) *AMSUA(REF)*.

Additional studies with AMV(REF)

Impact of MODIS AMVs

An experiment was run to evaluate the impact of the MODIS AMVs. This experiment is identical to AMV(REF) but with the MODIS AMVs removed. One can see that the impact of these AMVs on the 500 hPa geopotential height is very positive in the southern hemisphere (Figure 10(a)) and northern hemisphere (Figure 10(b)). The AMVs are clearly very important for observing the polar flow.

Comparison of AIRS with a combination of AMSU-A and AMSU-B

In the standard set of impact experiments using AMV(REF), Table 2, the AMV(REF)+AMSUA assimilation used all four AMSU-A instruments. This is a somewhat unfair comparison if one wants to directly compare the impact of AIRS data with that of AMSU-A. The AIRS channels used in the operational assimilation are mostly sensitive to temperature and moisture (McNally et al., 2004). A microwave assimilation experiment was therefore run adding one AMSU-A/B combination to the AMV(REF). This OSE was then compared with the AMV(REF)+AIRS assimilation referred to in Table 2. The results are shown in Figures 11(a) and 11(b).

In the southern hemisphere at day 2 the impact of AIRS and AMSU-A/B data on the geopotential height scores is similar but at days 5 and 7 the impact of the AIRS data becomes larger. However, the impact on tropical moisture, Figure 11(c), shows that AMSU-A/B data has a larger impact than AIRS in terms of the short-range forecast of tropical moisture.




Figure 10 (Above) Impact of MODIS AMVs on 500 hPa geopotential height for (a) southern hemisphere (20°–90°S) and (b) northern hemisphere (20°–90°N).




Figure 11 (Right) Comparison of AIRS with the two microwave instruments (AMSU-A and AMSU-B) on NOAA-16 for (a) 500 hPa geopotential height in the southern hemisphere, (b) 500 hPa geopotential height in the northern hemisphere and (c) 500 hPa relative humidity in the tropics.

Additional studies with AMSUA(REF)

AIRS channel combinations

For a number of reasons (including CPU time, memory size and file space constraints) the current operational system uses a reduced channel set for AIRS (and more recently IASI) radiances. The aim of this study was to explore which AIRS channels are the most important. Experiments have been run (summer only) by adding various AIRS channel combinations to the *AMSUA(REF)*, and also denying AIRS data from *CONTROL*. The results are shown in Figure 12 for 500 hPa geopotential height in the southern hemisphere. The respective performance of each scenario is consistent throughout the forecast range up to day 5.

When looking at the day 3 impact on the geopotential height at 500 hPa, one notices that:

- The positive impact of AIRS data can be seen from the differences between the data denial experiment and *CONTROL*. There is also a clear separation between *AMSUA(REF)*, and *AMSUA(REF)+AIRS*.
- *AMSUA(REF)+AIRS* is close to the *AIRS* experiment not using humidity channels, indicating that AIRS humidity channels do not have a large effect on the geopotential height at 500 hPa.
- The fact that *AMSUA(REF)+AIRS* is close to the experiment using only the 15 micron channels from AIRS suggests that these channels contribute most to the impact of this instrument.

Impact of the SSMI clear-sky and rain-affected radiances

In addition to the *AMSUA(REF)* plus all SSMI data (clear-sky and rain affected observations, *Bauer et al.*, 2002, 2006a and 2006b), *CONTROL* (all data) and *AMSUA(REF)* the following experiments have been performed to separate the effects of clear-sky and rain-affected SSMI radiances:

- *AMSUA(REF)* plus SSMI rain affected radiances.
- *AMSUA(REF)* plus SSMI clear-sky radiance radiances.
- *CONTROL* minus all SSMI (clear-sky and rain-affected) radiances.

Figure 13 shows the impact of denying and adding SSMI radiances (in the various configurations explained above) on the 850 and 500 hPa relative humidity forecast scores. The rain-affected radiances contribute more to the moisture forecast skill at 500 hPa whereas the clear-sky radiances are more important at 850 hPa. The experiment that combines both SSMI radiance types further improves the forecast. It is therefore clear that both types for SSMI radiances are important for the global moisture analysis.

Impact of AVHRR AMVs

During the study, a new experimental AMV product became available from CIMSS and it was decided to evaluate them in this OSE framework. These AMVs are produced from overlapping orbits from the AVHRR imager onboard the NOAA polar satellite series. Unfortunately this instrument does not have a water vapour channel like MODIS and this greatly reduces the amount of AMVs produced, particularly over the polar ice. The impact on mean scores is small but positive and can be best seen on the mean geographical rms forecast error difference with *AMSUA(REF)* for the 500 hPa geopotential height (Figure 14(a)). Very little impact is found in the northern hemisphere. For comparison a similar plot is shown when MODIS AMVs are assimilated instead (Figure 14(b)). The impact from the MODIS AMVs is similar to that of AVHRR AMVs over the Southern Ocean but extends further to the south over the polar ice.




Figure 12 Comparison of the impact of various AIRS channel combinations on 500 hPa geopotential height based on *AMSUA(REF)*.




Figure 13 Comparison of the impact of various SSMI usage configurations based on *AMSUA(REF)* for (a) 850 hPa relative humidity and (b) 500 hPa relative humidity.




Figure 14 Mean normalized 48-hour forecast error difference between (a) AVHRR AMVs and *AMSUA(REF)*, and (b) MODIS AMVs and *AMSUA(REF)* for the 500 hPa geopotential height.

Overall assessment and further prospects

All the space based sensors contribute in a positive way to the overall improvement of the ECMWF forecast system. Sensors like AMSU-A, AIRS and HIRS are clearly the most important for mass and wind forecasts. However the accuracy of the humidity forecast relies on AMSU-B/MHS, GEO CSRs and SSMI. The positive impact of AMVs (GEO and MODIS) and SCAT on the forecast is also clearly demonstrated.

At present, there are no plans to fly an instrument with MODIS-like water vapour channels on future polar satellites. This is a concern as the positive impact of MODIS AMVs in polar regions and mid-latitudes has clearly been demonstrated. The experiment using AVHRR derived AMVs show that their quality is similar to MODIS AMVs but the coverage is and will remain much poorer over the frozen regions due to a lack of water vapour channel on the instrument.

The studies also show that AIRS is the sensor that has the most impact on the mass field and experiments indicate that most of the impact comes from its 15 micron spectral band. Otherwise SSMI is vital for humidity analysis and the newly introduced cloud/rain affected SSMI radiances further improve the humidity analysis.

These experiments confirm the crucial impact of satellite data on the performance of the ECMWF NWP system. Since the completion of the OSEs, the importance of satellite data has further increased with, for example, the implementation of GPS radio-occultation observations or more recently the introduction of IASI. On the scientific side, further changes are expected in the near future that include the use of more infrared and microwave radiances in cloudy and rainy conditions, and an improved use of all types of satellite radiances over land and sea-ice.

Further Reading

Andersson, E., R. Dumelow, H. Huang, J.-N. Thépaut & A. Simmons, 2004: *Space/Terrestrial Link – Outline study proposal for consideration by EUCOS Management* (available from EUCOS Secretariat).

Andersson, E., J. Pailleux, J.-N. Thépaut, J.R. Eyre, A.P. McNally, G.A. Kelly & P. Courtier, 1994. Use of cloud-cleared radiances in three/four-dimensional variational data assimilation. *Q.J.R. Meteorol. Soc.*, **120**, 627–654.

Bauer, P., G. Kelly & E. Andersson, 2002: SSM/I radiance assimilation at ECMWF. In *Proc. ECMWF/GEWEX Workshop on Humidity Analysis*, Reading, United Kingdom, ECMWF, 167-175. Available from European Centre for Medium-Range Weather Forecasts, Shinfield Park, Reading RG2 9AX, UK.

Bauer, P., P. Lopez, A. Benedetti, D. Salmond & E. Moreau, 2006a: Implementation of 1D+4D-Var assimilation of precipitation affected microwave radiances at ECMWF, Part I: 1D-Var. *Q.J.R. Meteorol. Soc.*, **132**, 2277–2306.

Bauer, P., P. Lopez, A. Benedetti, D. Salmond, S. Saarinen & M. Bonazzola, 2006b: Implementation of 1D+4D-Var assimilation of precipitation affected microwave radiances at ECMWF, Part II: 4D-Var. *Q.J.R. Meteorol. Soc.*, **132**, 2307–2332.

Bouttier, F. & G. Kelly, 2001: Observing-system experiments in the ECMWF 4D-Var data assimilation system. *Q.J.R. Meteorol. Soc.*, **127**, 1469–1488.

Kelly, G. 1997: Influence of observations on the operational ECMWF system. *Bulletin of WMO*, **46**, 336–341.

Kelly, G., A. McNally, J.-N. Thépaut & M. Szyndel, 2004. Observing System Experiments of all main data types in the ECMWF operational system. In *Proc. of the Third WMO Workshop on the Impact of Various Observing Systems on NWP*, Alpbach, Austria, 9–12 March 2004.

McNally, A.P., P.D. Watts, J.A. Smith, R. Engelen, G. Kelly, J.-N. Thépaut & M. Matricardi, 2004: The assimilation of AIRS radiance data at ECMWF. In *Workshop on Assimilation of High Spectral Resolution Sounders in NWP*, 28 June-1 July 2004, Shinfield Park, Reading 73–88.

Thépaut, J.-N. & G.A. Kelly, 2007. Relative contributions from various terrestrial observing systems in the ECMWF NWP system. *Final Report EUCOS*, 23 June 2007 (available from EUCOS Secretariat).

© Copyright 2016

European Centre for Medium-Range Weather Forecasts, Shinfield Park, Reading, RG2 9AX, England

The content of this Newsletter article is available for use under a Creative Commons Attribution-Non-Commercial-No-Derivatives-4.0-ported Licence. See the terms at <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

The information within this publication is given in good faith and considered to be true, but ECMWF accepts no liability for error or omission or for loss or damage arising from its use.